


ID Clinical Update

Jonathan Vilasier Iralu, MD, MACP, FIDSA Indian Health Service Chief Clinical Consultant for Infectious Diseases



Virology:

- ✤ JN.1 is the most common variant
- No new concerns about vaccine and treatment efficacy





COVID Prevention: Spring vaccines

***** Special situation for people ages 65 years and older:

- People ages 65 years and older should receive 1 additional dose of any updated (2023–2024 Formula) COVID-19 vaccine at least 4 months following the previous dose of updated (2023–2024 Formula) COVID-19 vaccine.
- For initial vaccination with Novavax COVID-19 Vaccine, the 2-dose series should be completed before administration of the additional dose.



COVID Treatment Updates: January 2024

- The NIH COVID treatment guidelines are now available in the final version online.
 - <u>https://files.covid19treatmentguidelines.nih.gov/guidelines/covid19treatmentguidelines.pdf</u>

Thank you, NIH!



ID Guideline Updates

IDSA: No major updates

DHHS HIV antiretroviral guidelines:

- ♦ Age 40-75 years with 10 year ASCVD risk 5-20% → Initiate moderate intensity statin Rx (A1)
- Irrest risk < 5%, the panel recommends statin Rx (C1) : consider risk factors
- * Age < 40: lifestyle modifications are recommended</pre>

https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-arv/whats-new line adolescent-arv/whats-new line adolescent-adolescent-arv/whats-new line adolescent-arv/whats-new line adolescent-arv/whats-new line adolescent-arv/whats-new line adolescent-arv/whats-new line adolescent-arv/whats-new line adolescent-adolescent



ID Guideline Updates International AIDS Society (IAS-USA) ART update

- * Injectable Cabotegravir/Rilpivirine originally approved as a switch regimen for patients with viral suppression
- CAB-RPV now recommended for persons with viremia who have no other treatment options due to inability to take oral therapy
 - * Unable to take oral ART despite extensive efforts and clinical support
 - High risk for progression (CD4 <200 or AIDS-defining complications)</p>
 - Virus susceptible to CAB and RPV

https://jamanetwork.com/journals/jama/fullarticle/2816037



Indian Country Penicillin availability


*Penicillin is available from the prime vendor

*Keep on ordering and use it for

- Every patient
- Every encounter including clinic, hospital, and "home"
 Every stage



Indigi-IWTK is here!



Tests for:

- Syphilis
- HIV
- Gonorrhea
- Chlamydia
- Trichomonas
- HBV
- HCV

https://cih.jhu.edu/indigi-iwtk-i-want-the-kit/

Why use self collection kits?



Coming very soon...

Simple HIV PrEP guidelines



Being developed for you and your patients...

- HS STI Web page
 - Statistics
 - Treatment guidelines
 - *****Doxy-PEP
 - ♦ HIV PrEP
 - *Field Penicillin

Link to STI toolkit at NPTC Strategic Initiatives page

